

# Quick Review of Remote Sensing Basic Theory

Paolo Antonelli  
(Material Provided by Paul Menzel)


CIMSS  
University of Wisconsin-Madison

Benevento, June 2007



# Infrared (Emissive Bands)

Radiative Transfer Equation  
in the IR



# Relevant Material in Applications of Meteorological Satellites

## CHAPTER 2 - NATURE OF RADIATION

2.1	Remote Sensing of Radiation	2-1
2.2	Basic Units	2-1
2.3	Definitions of Radiation	2-2
2.5	Related Derivations	2-5


## CHAPTER 3 - ABSORPTION, EMISSION, REFLECTION, AND SCATTERING

3.1	Absorption and Emission	3-1
3.2	Conservation of Energy	3-1
3.3	Planetary Albedo	3-2
3.4	Selective Absorption and Emission	3-2
3.7	Summary of Interactions between Radiation and Matter	3-6
3.8	Beer's Law and Schwarzschild's Equation	3-7
3.9	Atmospheric Scattering	3-9
3.10	The Solar Spectrum	3-11
3.11	Composition of the Earth's Atmosphere	3-11
3.12	Atmospheric Absorption and Emission of Solar Radiation	3-11
3.13	Atmospheric Absorption and Emission of Thermal Radiation	3-12
3.14	Atmospheric Absorption Bands in the IR Spectrum	3-13
3.15	Atmospheric Absorption Bands in the Microwave Spectrum	3-14
3.16	Remote Sensing Regions	3-14

## CHAPTER 5 - THE RADIATIVE TRANSFER EQUATION (RTE)

5.1	Derivation of RTE	5-1
5.10	Microwave Form of RTE	5-28

# Energy conservation




$$\tau + a + r \neq 1$$


'ENERGY  
CONSERVATION'

$$\tau + a + r = 1$$


# Simple case with no atmosphere and opaque cloud



# Simple case with no atmosphere and semi-transparent cloud



# Radiative Transfer Equation




# Transmittance for Window Channels

$$\tau + a + r = 1$$

$\tau$  close to 1  
 $a$  close to 0


The molecular species in the atmosphere are not very active:

- most of the photons emitted by the surface make it to the Satellite
- if  $a$  is close to 0 in the atmosphere then  $\epsilon$  is close to 0, not much contribution from the atmospheric layers





# Transmittance for Absorption Channels




Absorption Channel:  
 $\tau$  close to 0  
 $a$  close to 1



One or more molecular species in the atmosphere is/are very active:

- most of the photons emitted by the surface will not make it to the Satellite (they will be absorbed)
- if  $a$  is close to 1 in the atmosphere then  $\epsilon$  is close to 1, most of the observed energy comes from one or more of the uppermost atmospheric layers


# Spectral Characteristics of Atmospheric Transmission and Sensing Systems




# Weighting Functions



**High resolution atmospheric absorption spectrum  
and comparative blackbody curves.**



# MODIS absorption bands



## Emission, Absorption

Blackbody radiation  $B_\lambda$  represents the upper limit to the amount of radiation that a real substance may emit at a given temperature for a given wavelength.

Emissivity  $\varepsilon_\lambda$  is defined as the fraction of emitted radiation  $R_\lambda$  to Blackbody radiation,

$$\varepsilon_\lambda = R_\lambda / B_\lambda .$$

In a medium at thermal equilibrium, what is absorbed is emitted (what goes in comes out) so

$$a_\lambda = \varepsilon_\lambda .$$

Thus, materials which are strong absorbers at a given wavelength are also strong emitters at that wavelength; similarly weak absorbers are weak emitters.

## Transmittance

Transmission through an absorbing medium for a given wavelength is governed by the number of intervening absorbing molecules (path length  $u$ ) and their absorbing power ( $k_\lambda$ ) at that wavelength. Beer's law indicates that transmittance decays exponentially with increasing path length

$$\tau_\lambda (z \rightarrow \infty) = e^{-k_\lambda u (z)}$$

where the path length is given by  $u (z) = \int_z^\infty \rho dz$ .

$k_\lambda u$  is a measure of the cumulative depletion that the beam of radiation has experienced as a result of its passage through the layer and is often called the optical depth  $\sigma_\lambda$ .

Realizing that the hydrostatic equation implies  $g \rho dz = -q dp$

where  $q$  is the mixing ratio and  $\rho$  is the density of the atmosphere, then

$$u (p) = \int_0^p q g^{-1} dp \quad \text{and} \quad \tau_\lambda (p \rightarrow 0) = e^{-k_\lambda u (p)}$$

## Emission, Absorption, Reflection, and Scattering

If  $a_\lambda$ ,  $r_\lambda$ , and  $\tau_\lambda$  represent the fractional absorption, reflectance, and transmittance, respectively, then conservation of energy says

$$a_\lambda + r_\lambda + \tau_\lambda = 1 .$$

For a blackbody  $a_\lambda = 1$ , it follows that  $r_\lambda = 0$  and  $\tau_\lambda = 0$  for blackbody radiation. Also, for a perfect window  $\tau_\lambda = 1$ ,  $a_\lambda = 0$  and  $r_\lambda = 0$ . For any opaque surface  $\tau_\lambda = 0$ , so radiation is either absorbed or reflected  $a_\lambda + r_\lambda = 1$ .

At any wavelength, strong reflectors are weak absorbers (i.e., snow at visible wavelengths), and weak reflectors are strong absorbers (i.e., asphalt at visible wavelengths).



## Radiative Transfer Equation

The radiance leaving the earth-atmosphere system sensed by a satellite borne radiometer is the sum of radiation emissions from the earth-surface and each atmospheric level that are transmitted to the top of the atmosphere. Considering the earth's surface to be a blackbody emitter (emissivity equal to unity), the upwelling radiance intensity,  $I_\lambda$ , for a cloudless atmosphere is given by the expression

$$I_\lambda = \varepsilon_\lambda^{\text{sfc}} B_\lambda(T_{\text{sfc}}) \tau_\lambda(\text{sfc} - \text{top}) + \sum_{\text{layers}} \varepsilon_\lambda^{\text{layer}} B_\lambda(T_{\text{layer}}) \tau_\lambda(\text{layer} - \text{top})$$

where the first term is the surface contribution and the second term is the atmospheric contribution to the radiance to space.

When reflection from the earth surface is also considered, the Radiative Transfer Equation for infrared radiation can be written

$$I_{\lambda} = \varepsilon_{\lambda}^{\text{sfc}} B_{\lambda}(T_s) \tau_{\lambda}(p_s) + \int_{p_s}^0 B_{\lambda}(T(p)) F_{\lambda}(p) [d\tau_{\lambda}(p) / dp] dp$$

where


$$F_{\lambda}(p) = \{ 1 + (1 - \varepsilon_{\lambda}) [\tau_{\lambda}(p_s) / \tau_{\lambda}(p)]^2 \}$$

The first term is the spectral radiance emitted by the surface and attenuated by the atmosphere, often called the boundary term and the second term is the spectral radiance emitted to space by the atmosphere directly or by reflection from the earth surface.


The atmospheric contribution is the weighted sum of the Planck radiance contribution from each layer, where the weighting function is  $[d\tau_{\lambda}(p) / dp]$ . This weighting function is an indication of where in the atmosphere the majority of the radiation for a given spectral band comes from.

# Earth emitted spectra overlaid on Planck function envelopes


High resolution atmospheric absorption spectrum and comparative blackbody curves.



# Weighting Functions



CO2, strat temp  
 CO2, strat temp  
 CO2, upper trop temp  
 CO2, mid trop temp  
 CO2, lower trop temp  
 H2O, lower trop moisture  
 H2O, dirty window



window  
 O3, strat ozone  
 H2O, lower mid trop moisture  
 H2O, mid trop moisture  
 H2O, upper trop moisture

# Conclusion

- Radiative Transfer Equation (IR): models the propagation of terrestrial emitted energy through the atmosphere